

Relationship between Procyanidin and Flavor Contents of Cocoa Liquors from Different Origins

CHRISTINE COUNET, CAROLINE OUWERX, DELPHINE ROSOUX, AND SONIA COLLIN*

Unité de Brasserie et des Industries Alimentaires, Université catholique de Louvain,
 Croix du Sud, 2/Bte7, B-1348 Louvain-la-Neuve, Belgium

The flavor of eight cocoa liquors of different origins (Africa, America, and Asia) and different varieties (Fine grades: *criollo*, *trinitario*, and *nacional*. Bulk-basic grade: *forastero*.) was analyzed by headspace solid-phase microextraction mass spectrometry (HS-SPME-MS). Their procyanidin contents were quantified by HPLC–UV (280 nm). Fine varieties with short fermentation processes proved to contain more procyanidins, while *criollo* from New Guinea and *forastero* beans showed the highest aroma levels. The levels of cocoa aroma compounds formed during roasting are shown to vary directly with bean fermentation time and inversely with residual procyanidin content in cocoa liquor. Measurement of antioxidant activity in cocoa liquor proved to be a useful tool for assessing residual polyphenols.

KEYWORDS: Cocoa origins; chocolate; procyanidins; flavor; AAPH antioxidant activity

INTRODUCTION

The flavor and polyphenol contents of cocoa liquors depend on cocoa bean variety (genotype), postharvest processes (fermentation and drying), and roasting conditions (1–4). Mainly three varieties of cocoa beans are produced worldwide: *forastero* (bulk grade, 70% of the world production), *criollo* (fine grade), and their hybrid, *trinitario* (fine grade) (5). Cacao from Ecuador (Arriba) is viewed as a third fine variety: *nacional* (6). Bulk cocoas usually exhibit strong, harsh flavors while fine cocoas are perceived as more aromatic or smoother (4). Cocoa bean fermentation and drying contribute to developing cocoa–chocolate flavors by increasing levels of amino acids and sugars, reagents of the Maillard reactions that occur during roasting. Through these postharvest processes, polyphenols are usually lost by diffusion, browning, and oxidative polymerization.

Dried fermented cocoa beans from Southeast Asia and the South Pacific are characterized by a higher acidity (higher levels of lactic and acetic acids) than those of South African origin. Of the four acids analyzed (lactic, acetic, citric, and oxalic), only oxalic acid seems to have a positive impact on chocolate taste (7).

Each cocoa liquor exhibits its own organoleptic properties. For instance, Cameroon liquors are famous for their bitterness while Ecuador liquors are known for their raisin–fruity flavor. Bailey et al. (1) detected only quantitative variations in aroma between origins of dried fermented cocoa beans (these authors investigated the Accra, Arriba, Bahia, Sanchez, and Trinidad origins). According to Ziegler (8), fine dried fermented cocoa beans (Ecuador, Trinidad, and Venezuela) contain up to 8 times more linalool than bulk-basic ones (Ghana, Ivory Coast, Brazil). Linalool might thus be responsible for the flowery–tea note

found in fine varieties. In addition, Muggler-Chavan et al. (9) found higher butanol, isopentanol, hexanol, and octanol levels in *criollo*–*trinitario*–*nacional* dried fermented cocoa beans (Puerto Cabello, Arriba, Trinidad, and Caracas) than in four *forastero* origins (Carupano, Bahia, Para, and Accra). On the other hand, high concentrations of phenol, guaiacol, 2-phenylbutenal, and γ -butyrolactone characterize the Bahia beans known for their typical smoked ham odor. Also noteworthy are the higher levels of 2-methylpropanal and 3-methylbutanal in Caracas and Trinidad dried fermented beans (10). As far as other Maillard products are concerned, Reineccius et al. (11) report that roasting leads to higher levels of pyrazines in well-fermented beans (Ghana, Bahia) than in less-fermented (Arriba) or unfermented (Sanchez, Tabasco) materials.

Owing to their lower astringency and bitterness, mainly imparted by polyphenols according to refs 3 and 12, *criollo* beans are often less fermented than the *forastero* varieties. The *N* index, defined as the soluble nitrogen percentage [$(N_{\text{soluble}}/N_{\text{total}}) \times 100$] is used to assess the intensity of the fermentation process. On a scale of shorter to longer fermentation times, the order is usually as follows: Machala (Ecuador) < Sanchez (Dominican Republic) < Para (Brazil) < Costa Rica < Carupano (Venezuela) < Indonesia < Grenada < Jamaica < Fernando Po (Republic of Ecuador/Guinea) < Puerto Cabello (Venezuela) < Trinidad < Arriba (Ecuador) < Ivory Coast < Accra (Ghana) < New Guinea < Cameroon < Lagos (Nigeria) < San Thome < Bahia (Brazil) (13).

As recently shown by Counet et al. (14), procyanidins, the main class of polyphenols in cocoa products (Figure 1), impart not only astringency and bitterness to chocolate but also its exceptional antioxidant activity. The aim of the present work was therefore to investigate how the procyanidin level of cocoa liquors of different origins might be predicted (14). Eight different samples were analyzed by HS-SPME-GC-MS for their

* To whom correspondence should be addressed. Telephone: (+ 32) 10 47 29 13. Fax: (+ 32) 10 47 21 78. E-mail: collin@inbr.ucl.ac.be.




Figure 1. Example of procyanidin monomers and dimers found in cocoa products.

aroma content and by HPLC–UV/MS so as to quantify procyanidin in aqueous acetone extracts of defatted cocoa liquors. Measurement of antioxidant activity was also performed by using the AAPH [2,2'-azobis(2-amidinopropane)dihydrochloride]-induced linoleic oxidation assay.

EXPERIMENTAL PROCEDURES

Materials. Eight cocoa liquors from different origins and varieties were supplied by Belcolade (Eremodegem, Belgium): Ivory Coast and Ghana (*forastero*), New Guinea, Java, Venezuela and Madagascar (*criollo*), Ecuador (*nacional*), Trinidad (*trinitario*).

Chemicals. Acetone (99.9%), (-)-epicatechin, (+)-catechin, and most flavor compounds were from Sigma-Aldrich (Bornem, Belgium). Only 1-methylpropyl acetate, 2-pentanol acetate, 3-methyl-1-butanol acetate, and 2-methyl-1-butanol acetate were synthesized from the corresponding alcohol and acids in the presence of sulfuric acid. Methanol (99.9%) and dichloromethane (99.9%) were purchased from Romil (Cambridge, U.K.). Acetic acid (99.8%) was from Acros (Geel,

Belgium), and diethyl ether (99.5%) from Fluka (Buchs, Switzerland). Aqueous solutions were made with Milli-Q (Millipore, Bedford, MA) double-distilled water (resistance = 18 mΩ).

Procyanidin Analysis of Cocoa Liquors (14). *Lipid Removal.* Cocoa liquor (70 g) was reduced to a powder with a mixer and introduced into the Soxhlet extractor (Waterkeyn, Belgium) by using a filtration cartridge (Schleicher & Schüll, Germany). Lipids were removed for 24 h with diethyl ether (375 mL). Defatted cocoa liquor (50 g) was finally obtained.

Procyanidin Extraction. Defatted cocoa liquor (10 g) was extracted three times with 50 mL of solvent (3 × 1 h, 25 °C). The organic solvent used for procyanidin extraction was acetone mixed with water and acetic acid (70:28:2, v/v). After each extraction, the suspension was centrifuged for 10 min at 3000g, and the supernatant was collected. After filtration to remove residual particles, the combined supernatants were concentrated by rotary evaporation under partial vacuum (40 °C) to obtain about 50 mL of extract.

Procyanidin Extracts Purification. A 10 g C18 Sep-Pack cartridge (Waters, Millipore) was preconditioned with methanol and then with deionized water. About 50 mL of a procyanidin mixture was loaded on the cartridge, and sugars were removed with 200 mL of deionized water. Procyanidins were then eluted with 30 mL of acetone–water–acetic acid (70:28:2, v/v). The eluates were concentrated by rotary evaporation under partial vacuum (40 °C) and freeze-dried.

High Performance Liquid Chromatography Analysis of Procyanidin Extracts (HPLC–UV). A SpectraSystem (Finnigan Mat, San Jose, CA) equipped with a SCM degasser, an AS3000 autosampler, a P4000 quaternary pump, and a diode array detector UV6000LP at 280 nm was used for quantification (identification previously checked by mass spectrometry according to ref 14). The system was controlled with Xcalibur software version 1.2 (Finnigan Mat). Procyanidins were separated on a Phenomenex 5 μm normal-phase Luna silica column, 250 × 4.6 mm i.d. (Bester, Holland) at 25 °C. Separations were carried out at a flow rate of 1 mL/min with a linear gradient from A (dichloromethane) to B (methanol) and a constant 4% level of C (acetic acid and water, 1:1, v/v). Gradient elution was 14–28% B, 0–30 min; 28–50% B, 30–60 min; 50–86% B, 60–65 min; 65–70 min isocratic. Procyanidin extract (10 mg) was diluted in 1 mL of methanol before injections in duplicate with the 20 μL Rheodyne loop (Berkeley, CA).

Antioxidant Assay: AAPH Method. The reduction power of cocoa liquor extracts was measured by a method developed in our laboratory by Liégeois et al. (15). The oxidation of linoleic acid was induced by 2,2'-azobis(2-amidinopropane)dihydrochloride (AAPH) in an aqueous




Figure 2. HPLC–UV (280 nm) chromatogram of the Madagascar cocoa liquor procyanidin extract (P1 to P8 = monomeric to octameric procyanidins).




Figure 3. Total (a and b) and detailed (c) concentrations of procyanidins (P1 to P8 = monomeric to octameric procyanidins) in extracts of cocoa liquors of different origins (extraction and HPLC–UV injection each in duplicate, CV = 2.8–1.7%): (a) mg/100 g of dry extract; (b and c) mg/kg of cocoa liquor (recovery factor set at 100%). (C) = *criollo*, (T) = *trinitario*, (F) = *forastero*, and (N) = *nacional*.

dispersion in the absence or presence of antioxidant. The rate of oxidation at 37 °C was monitored by recording the increase in absorption at 234 nm caused by conjugated diene hydroperoxides. A Shimadzu (Antwerp, Belgium) UV–visible 240 spectrophotometer equipped with an automatic sample positioner allowed analysis of six samples per minute. In all cases, the measurements were run in duplicate against the buffer and compared with the case of a separate AAPH-free control to check for any spontaneous oxidation.

Flavor Analysis of Cocoa Liquors. *Headspace Solid-Phase Microextraction (HS-SPME).* Twenty-three millimeters of PDMS/Carboxen fiber (Supelco, Bellefonte, PA) was introduced into the headspace of vials (Chromacol Ltd, Lokeren, Belgium) containing 2 g of cocoa

liquor and equipped with a magnetic CrimpCap (20 mm in diameter) and a silicon–Teflon septum (Interscience, Louvain-la-Neuve, Belgium). The volatile compounds of 2 g of cocoa liquor were adsorbed onto the fiber for 20 min at 25 °C. They were then automatically desorbed for 5 min at 250 °C in the gas chromatography injector (penetration of the fiber: 54 mm).

Gas Chromatography–Mass Spectrometry Analysis (GC-MS) (18). The gas chromatograph was a Trace GC (Finnigan Mat, San Jose, CA) equipped with a split–splitless injector and an MS detector linked to a computer with Xcalibur software version 1.2 (Finnigan Mat). Compounds were separated using a 50 m × 0.32 mm i.d., wall coated open tubular (WCOT) nonpolar CP-Sil5-CB capillary column (film

Table 1. Compounds Tentatively Identified by SPME-GC-MS-RI (Retention Index) in the New Guinea Cocoa Liqueur

T_r (min)	RI	compound	T_r (min)	RI	compound
6.01	665	3-methylbutanal ^{a,b,d,f}	18.31	928	5-methyl-2-furancarboxaldehyde ^{a,c,e}
6.12	668	2-methylbutanal ^{a,f}	18.63	933	benzaldehyde ^{a,c,e,f}
6.71	686	pentanal ^f	20.16	949	dimethyl trisulfide ^{a,b,d}
6.82	689	2-pentanol ^f	21.82	973	2-ethyl-6-methylpyrazine ^{a,e,f}
8.26	732	dimethyl disulfide ^a	22.23	978	trimethylpyrazine ^{a,b}
8.46	738	1-methylpropyl acetate	22.62	982	myrcene ^c
8.95	752	toluene ^a	24.54	1005	benzyl alcohol ^{a,e,f}
9.71	775	dihydro-2-methyl-3(2H)-furanone ^{a,e} and hexanal ^{d,f}	26.14	1021	2-acetylpyrrole ^{a,c,e}
10.67	802	methylpyrazine ^{a,c}	26.66	1026	limonene ^c
10.91	806	3-methylbutanoic acid ^{b,d}	27.82	1037	acetophenone ^{c,e,f}
11.39	815	2-methylbutanoic acid ^{b,d}	29.75	1056	3(or 2)-ethyl-2(or 3),5-dimethylpyrazine ^{a-f}
12.09	828	2-pentanol acetate ^c	30.39	1062	tetramethylpyrazine ^{a,c,e}
13.46	853	3-methyl-1-butanol acetate	31.20	1070	2-nonanone ^{c,f}
13.61	856	2-methyl-1-butanol acetate	31.72	1074	oxyde de linalool ^a
14.13	865	2-heptanone ^{a,e,f}	32.78	1084	β -phenylethanol ^{a,b,c,e}
14.95	880	2-heptanol ^{a,c}	33.80	1094	2-isopropyl-5-methylhex-2-enal
15.32	887	2,5-dimethylpyrazine ^{a,c,e}	38.82	1138	2,3,5-trimethyl-6-ethylpyrazine or 3,5-diethyl-2-methylpyrazine ^{a,e}
15.64	893	ethylpyrazine ^{a,c,e}	43.62	1178	ethyloctanoate ^{a,c,f}
15.86	897	2,3-dimethylpyrazine ^{a,c,e} and 3-hydroxy-2-butanone (acetoin) ^a	49.43	1226	2-phenylethyl acetate ^{a,b,c,d}
16.25	903	4-hydroxy-2-butanone	50.84	1237	α -ethylidene benzene acetaldehyde (2-phenylbut-2-enal) ^{a,e}
			74.07	1485	5-methyl-2-phenyl-2-hexanal ^{a,c,e}

T_r = retention time (min), RI = retention index (min). ^a Compound previously identified in a dark chocolate by (18). ^b Compound previously reported as key odorant in bitter chocolate by (19). ^c Compound previously identified in a dark semi-sweet chocolate before conching by (23). ^d Compound previously reported as key odorant in milk chocolate by (20). ^e Compound previously identified in milk chocolate by (24). ^f Compound previously identified in chocolate (type not defined) by (25).




Figure 4. Volatile compounds potential (scored as the sum of all peak areas) of each cocoa liqueur. (C) = *criollo*, (T) = *trinitario*, (F) = *forastero*, and (N) = *nacional*.

thickness: 1.2 μm) (Varian, Sint-Katelijne-Waver, Belgium). The oven temperature was programmed from 36 to 85 $^{\circ}\text{C}$ at 20 $^{\circ}\text{C}/\text{min}$, to 145 $^{\circ}\text{C}$ at 1 $^{\circ}\text{C}/\text{min}$, and to 250 $^{\circ}\text{C}$ at 3 $^{\circ}\text{C}/\text{min}$ and then was held constant at 250 $^{\circ}\text{C}$ for 30 min. The injection temperature, the split flow, and the splitless time were respectively 250 $^{\circ}\text{C}$, 20 mL/min, and 1 min. The retention index (RI) was calculated by using *n*-alkanes as references. MS analyses were carried out with a Trace MS quadrupole mass spectrometer (Finnigan Mat, San Jose, CA). Electron impact mass spectra were recorded at 70 eV (2.45 scan per second) with a 40–400 amu range.

Flavor Quantification. Before analysis, increasing concentrations of commercial standards (aromas) were added to the Venezuelan cocoa liqueur. The standard addition curves obtained in this way were then used to quantify compounds in other cocoa samples.

RESULTS AND DISCUSSION

Procyanidins in Cocoa Liqueur Extracts. As depicted in Figure 2 for the Madagascar origin, procyanidin monomers

(P1), dimers (P2), and so forth up to octamers (P8) were found in our eight cocoa liqueur extracts (HPLC–UV data compared with MS results previously reported for chocolate extracts (14)). Yet significant differences (as much as 8-fold) in total procyanidin concentration were observed between the origins, with the order being Madagascar > Java > Trinidad > Ecuador > Venezuela > Ivory Coast > Ghana > New Guinea (Figure 3a). Most of the fine-cocoa liqueur extracts (less fermented) emerged at the top of the range [Madagascar (*criollo*), Java (*criollo*), Trinidad (*trinitario*), and Ecuador (*nacional*)] while bulk cocoa liquors from Africa [Ivory Coast and Ghana (*forastero*), highly fermented] proved very poor in such compounds. Although it belongs to the *criollo* variety, the New Guinea extract ranked lowest on our scale. This is probably also due to the long fermentation applied.

Taking into account the yield of each cocoa liqueur, milligrams of procyanidins per 100 g of dry extract were converted into




Figure 5. Details of the HS-SPME-MS chromatograms between 9 and 13 min (identical attenuation in all cases): (1) hexanal and dihydro-2-methyl-3(2*H*)-furanone (the latter is found in only long-fermentation cocoa liquors); (2) methylpyrazine; (3) 3-methylbutanoic acid; (4) 2-methylbutanoic acid; (5) 2-pentanol acetate.

milligrams per kilogram of cocoa liquor (extraction yield set at 100%; **Figure 3b**). The total amount of procyanidins calculated in this way ranged between 2200 and 8300 ppm (0.22–0.83%). This means a difference as high as 6100 ppm between origins. The richer origin (Madagascar) was found to contain at least 3524 ppm P1, 1522 ppm P2, 1236 ppm P3, 821 ppm P4, 513 ppm P5, 326 ppm P6, 169 ppm P7, and 118 ppm P8 (**Figure 3c**). Because of the current unavailability of procyanidin oligomer standards, no standard addition was applied except for (–)-epicatechin (100% recovery factor). Hence, the lower levels measured for P4 to P8 might be partially due to less efficient recovery after extraction.

Flavor Analysis of Cocoa Liquors. SPME-GC-MS was used to identify 43 compounds in each cocoa liquor studied. The main compounds investigated were alcohols, esters, aldehydes,




Figure 6. Concentrations of five flavor compounds synthesized during the roasting step in different cocoa liquors.

ketones, hydrocarbons, nitrogen and oxygen heterocycles, nitriles, and sulfides (**Table 1**). Among them, five had never been described before in chocolate: 1-methylpropyl acetate, 3-methyl-1-butanol acetate, 2-methyl-1-butanol acetate, 4-hydroxy-2-butanone, and 2-isopropyl-5-methylhex-2-enal.

According to their volatile compounds potential (scored as the sum of all peak areas), cocoa liquors should be classified as follows: New Guinea \gg Ivory Coast \approx Ghana \gg Venezuela, Java, Trinidad, Ecuador, Madagascar (**Figure 4**). Fine cocoa liquors emerged as much less rich than the bulk-basic ones,




Figure 7. Correlation between the procyanidin level and (a) the 2,5-dimethylpyrazine content and (b) the total level of five compounds synthesized during roasting. (C) = *criollo*, (T) = *trinitario*, (F) = *forastero*, and (N) = *nacional*.

despite their higher quality according to consumer preferences (16). The chemical nature of the most potent flavors and how they are balanced by others (1, 17) are probably of prime importance in allowing perception of the fine fruity profile characterizing *criollo* varieties (16).

An enlargement of the 9–13 min zone of the chromatograms is depicted in Figure 5. The well-fermented cocoa liquors from New Guinea, Ghana, and Ivory Coast gave rise to more numerous and intense peaks than the fine varieties. Especially dihydro-2-methyl-3-(2*H*)-furanone (no. 1) and methylpyrazine (no. 2) proved specific to these three polyphenol-low liquors.

Five flavors synthesized by Maillard reactions during the roasting step (18–20) were quantified in each cocoa liquor: methylpyrazine (hazelnut-green), 2,5-dimethylpyrazine, trimethylpyrazine (cocoa-roasted-green), 3-ethyl-2,5-dimethylpyrazine (roasted-smoky-praline), and dimethyl disulfide (derived from methional and recognized as exhibiting a cocoa-like odor in synergy with 3-methylbutanal (21)) (see Figure 6). Bulk varieties from Africa and the New Guinea sample contained higher amounts of all these compounds, most probably due to higher levels of precursors synthesized through fermentation.

Also to be emphasized in the SPME data are the huge amounts of tetramethylpyrazine (590 ppm) and α -phenylethanol (512 ppm), respectively, in the cocoa liquors from Java and Venezuela. The former might impart the typical flowery–lemon note usually perceived with the Venezuelan cocoa liquor.

Correlation between the Procyanidin Level and Flavor Contents of Different Cocoa Liquors. To predict more easily the procyanidin content ([PC]) of a cocoa liquor, we tried to establish a mathematical correlation with its flavor content. As depicted in Figure 7a, a negative correlation with 2,5-dimethylpyrazine was obtained for four fine varieties. Only *nacional*




Figure 8. Correlation between the antioxidant activity (expressed as the inhibition time resulting from the presence of cocoa liquor extract) and the procyanidin level of different cocoa liquors. (C) = *criollo*, (T) = *trinitario*, (F) = *forastero*, and (N) = *nacional*.

cocoa liquor from Ecuador appears very different, but literature indicates that it is an independent variety from a genotype point of view (22). In the three long-fermentation origins, the best indicators of P1 to P8 procyanidin content proved to be the levels of five major compounds synthesized during roasting (Figure 7b). A large number of samples should of course be analyzed to confirm these preliminary results.

Correlation between the Procyanidin Level and Antioxidant Activities of Different Cocoa Liquors. As Figure 8 shows, quick measurement of the inhibition time observed in the AAPH-induced (2,2'-azobis(2-amidinopropane) dihydrochloride) oxidation assay after addition of cocoa liquor extract is another interesting way to predict the procyanidin level. The higher the antioxidant activity, the higher the level of polyphenols, suggesting that procyanidins are the best radical scavengers in cocoa liquors, despite the presence of high amounts of melanoidins (14).

CONCLUSIONS

The fermentation time of cocoa beans seems to be the key factor controlling the synthesis of aromas such as methylpyrazine, 2,5-dimethylpyrazine, trimethylpyrazine, 3-ethyl-2,5-dimethylpyrazine, and dimethyl disulfide. Expectedly, therefore, a negative correlation between procyanidins and some of these compounds emerged from our data. The antioxidant activity measured as the inhibition time found in the AAPH-induced linoleic oxidation assay appears as an easy way to predict the procyanidin level in chocolate factories.

ACKNOWLEDGMENT

The authors are grateful to Ir. C. Ouwere (Belcolade) and Dr. F. Arnaut (Puratos Group) for supplying cocoa liquor samples.

LITERATURE CITED

- Bailey, S.; Mitchell, D.; Bazinet, M.; Weurman, C. Studies of the volatile components of different varieties of cocoa beans. *J. Food Sci.* **1962**, *27*, 165–170.
- Bainbridge, J.; Davies, S. The essential oil of cocoa. *J. Chem. Soc.* **1912**, *101*, 2209–2221.
- Clapperton, J.; Yow, S.; Chan, J.; Lim, D.; Lockwood, R.; Romanczyk, L.; Hammerstone, J. The contribution of genotype to cocoa (*Theobroma cacao* L) flavour. *Trop. Agric.* **1994**, *71*, 303–308.
- Kattenberg, H.; Kemming, A. The flavor of cocoa in relation to the origin and processing of the cocoa beans. In *Food Flavors, Ingredients and Composition*; Charalambous, G. Ed.; Elsevier Science: Amsterdam, The Netherlands, 1993; pp 1–22.

- (5) Hoskin, J. C. Sensory properties of chocolate and their development. *Am J. Clin. Nutr.* **1994**, *60*, 1068S–1070S.
- (6) Despréaux, D. Le cacaoyer et la cacao-culture. In *Cacao et chocolat—production, utilisation, caractéristiques*; Pontillon, J., Ed.; Technique & Documentation Lavoisier: Paris, France, 1998; pp 44–93.
- (7) Holm, C.; Aston, J.; Douglas, K. The effects of the organic acids in cocoa on the flavour of chocolate. *J. Sci. Food Agric.* **1993**, *61*, 65–71.
- (8) Ziegleder, G. Linalool contents as characteristic of some flavour grade cocoas. *Lebensm. Unters. Forsch.* **1990**, *191*, 306–309.
- (9) Muggler-Chavan, F.; Reymond, D. Constituants aromatiques de cacao de diverses provenance. *Trav. Chim. Alim. Hyg.* **1967**, *58*, 466–473.
- (10) Hoskin, J.; Dimick, P. Chemistry of flavour development in chocolate. In *Industrial chocolate manufacture and use*; Beckett, S. T., Ed.; Blackie Academic & Professional: New York, 1994.
- (11) Reineccius, G.; Keeney, P.; Weissberger, W. Factors affecting the concentration of pyrazines in cocoa beans. *J. Agric. Food Chem.* **1972**, *20*, 202–206.
- (12) Luna, F.; Crouzillat, D.; Cirou, L.; Bucheli, P. Chemical composition and flavor of ecuadorian cocoa liquor. *J. Agric. Food Chem.* **2002**, *50*, 3527–3532.
- (13) Rohan, T. A.; Stewart, T. The precursors of chocolate aroma: production of free amino acids during fermentation of cocoa beans. *J. Food Sci.* **1967**, *32*, 395–398.
- (14) Counet, C.; Collin, S. Effect of the number of flavanol units on the antioxidant activity of procyanidin fractions isolated from chocolate. *J. Agric. Food Chem.* **2003**, *51*, 6816–6822.
- (15) Liégeois, C.; Lermusieau, G.; Collin, S. Measuring antioxidant efficiency of wort, malt, and hops against the 2,2'-azobis(2-amidinopropane)dihydrochloride-induced oxidation of an aqueous dispersion of linoleic acid. *J. Agric. Food Chem.* **2000**, *48*, 1129–1134.
- (16) Jinap, S.; Dimick, P. S.; Hollender, R. Flavour evaluation of chocolate formulated from cocoa beans from different countries. *Food Control* **1995**, *6*, 105–110.
- (17) Baigrie, B. Cocoa flavour. In *Understanding natural flavor*; Piggott, J. R., Paterson, A., Eds.; Blackie Academic & Professional—Chapman & Hall: Glasgow, U.K., 1994; pp 268–282.
- (18) Counet, C.; Callemien, D.; Ouwerx, C.; Collin, S. Use of GC-olfactometry to identify key odorant compounds in dark chocolate. Comparison of samples before and after conching. *J. Agric. Food Chem.* **2002**, *50*, 2385–2392.
- (19) Schieberle, P.; Pfner, P. Characterization of key odorants in chocolate. In *Flavor Chemistry: 30 Years of Progress*; Teranishi et al., Eds.; Kluwer Academic/Plenum: New York, 1999; pp 147–153.
- (20) Schnermann, P.; Schieberle, P. Evaluation of key odorants in milk chocolate and cocoa mass by aroma extract dilution analyses. *J. Agric. Food Chem.* **1997**, *45*, 867–872.
- (21) Lopez, A.; Quesnel, V. The contribution of sulphur compounds to chocolate aroma. In *Proceedings of the 1st International Congress on Cocoa and Chocolate Research*; Inst. Lebensmitteltechnologie Verpack: Munich, Germany, 1974; pp 92–104.
- (22) Lerceteau, E. Diversité génétique, recherche de QTL et analyse des profils protéiques des fèves de *Theobroma cacao L.* pendant la fermentation. Thèse de doctorat, Université d'Orsay, Paris XI, France, 1996.
- (23) Manière, D.; Dimick, P. Effects of conching on the flavor and volatile components of dark semi-sweet chocolate. *Lebensm. Wiss. Technol.* **1979**, *12*, 102–107.
- (24) Ziegleder, G.; Stojavic, E. Lagerungsbedingte veränderungen im aroma von milchschokoladen. *Z. Lebensm. Unters. Forsch.* **1988**, *186*, 134–138.
- (25) Ghizzoni, C.; Del Popolo, F.; Colombo, E.; Poretta, S. Composition of volatile fraction of industrial chocolate. *Ital. Food Beverage Technol.* **1995**, *5*, 3–13.

Received for review March 3, 2004. Accepted June 2, 2004. The authors are grateful for the “Grant F.N.R.S. – Spadel” and to the F.N.R.S. for financial support.

JF040105B